

A comment on Finite Temperature Correlations in Integrable QFT

H. Saleur

Department of Physics
University of Southern California
Los Angeles, CA 90089-0484

I discuss and extend the recent proposal of Leclair and Mussardo for finite temperature correlation functions in integrable QFTs. I give further justification for its validity in the case of one point functions of conserved quantities. I also argue that the proposal is not correct for two (and higher) point functions, and give some counterexamples to justify that claim.

In a recent paper [1], A. Leclair and G. Mussardo have come up with a very interesting suggestion to compute correlation functions in integrable quantum field theories at finite temperature. In a nutshell, what they propose is to identify the usual particle and hole excitations over the “thermal ground state” [2], with their non trivial dressed energies and momenta (as obtained by solving the thermodynamic Bethe ansatz equations), but still take, for the physical operators, the undressed form factors.

The proposal in [1] is quite exciting, as finite temperature correlation functions are quantities of the highest interest (in particular for comparison with experiments). However, it is fair to say that no real demonstration is provided in [1] except for some one point functions, and that the formula in [1] are merely a tentatizing guess.

I will argue in this note that, while the formula proposed in [1] for one point functions are, at least for some operators (the densities of conserved quantities), correct, the ones for two and higher point functions generally do not hold. The reason for this is quite simple and physical: the formula proposed in [1] does not take into account the effects of the dressing in the form-factors. This is fine for one point functions: since the *same* multiparticle states are on the left and on the right side of the correlator, there is no motion of the thermal ground state [2] due to interactions, and the formula for bare form factors can still be used. In contrast, for two point functions, one has to consider intermediate states where particle or holes have been created; this leads to a displacement of the thermal ground state, and a dressing of the form-factors, that has to be taken into account - there is just no reason for the bare form factors to still be relevant there.

One of the difficulties in assessing the validity of [1] is that very few finite temperature correlators are known exactly in integrable theories, and that the expressions proposed in [1], even if conceptually quite simple, are nevertheless very hard to compute explicitly ¹. However, as I will argue below, the argument presented in [1] is quite general, and holds just as well for correlators evaluated in other thermodynamic ensembles. The case of a chemical potential at $T = 0$ turns out to be particularly simple, and will allow me to put the two point functions of [1] to a serious test, that they unfortunately fail.

Let me now proceed and discuss one point functions first. I will only consider operators $O(x)$ (I refer to them, a bit incorrectly, as conserved quantities) for which the quantity $O = \int O(x)dx$ acts diagonally on multiparticle states, with one particle eigenvalues $o(\theta)$.

¹ A possibility might be to investigate the low temperature limit, and compare it with results obtained in [3], but I won't do this here.

Examples of this include the energy and momentum operators, and after a very slight modification, the trace of the stress energy tensor considered in [1]. By translational invariance, it thus follows that

$$\frac{\langle \theta_n, \dots, \theta_1 | O(x) | \theta_1, \dots, \theta_n \rangle}{\langle \theta_n, \dots, \theta_1 | \theta_1, \dots, \theta_n \rangle} = \frac{1}{L} \sum_{i=1}^n o(\theta_i),$$

L the system length.

I want now to compute the average of O at finite temperature, and in the presence of possible other thermodynamic couplings. I will restrict for notational simplicity to a theory with a single particle, and consider, in addition to the temperature, the presence of a chemical potential. Generalizations are quite straightforward. Due to the non trivial S matrix of the theory, the density of allowed states P ($P = \rho + \rho^h$) obeys (I set $\hbar = 1$ here):

$$2\pi P(\theta) = Lm \cosh \theta + 2\pi \Phi \star \rho(\theta), \quad (1)$$

where I defined $f \star g(\theta) = \int f(\theta - \theta') g(\theta') \frac{d\theta'}{2\pi}$. In (1), the kernel $\Phi = \frac{1}{2i\pi} \frac{d}{d\theta} \ln S$, where S is the scattering matrix.

The average of O is computed using the thermodynamic Bethe ansatz method: configurations are weighed with a Boltzmann weight $\exp \left[-\frac{E - TS - \mu N}{T} \right]$ where T is the temperature, S the entropy, μ the chemical potential, and N the number of particles, and saddle point equations are written, leading to the TBA equations. Introducing the pseudo energy ϵ defined by $\frac{\rho^h}{\rho} = e^\epsilon$, the TBA equation at temperature T ($T = \frac{1}{R}$) read

$$\epsilon(\theta) = mR \cosh \theta - \Phi \star \ln (1 + e^{-\epsilon + \mu R}), \quad (2)$$

It can be shown with the usual argument [4] that $2\pi P = L \left. \frac{\partial \epsilon}{\partial R} \right|_{\mu R \text{ fixed}}$. From this the average of O follows:

$$\begin{aligned} \langle O(x) \rangle &= \frac{1}{L} \int o(\theta) \rho(\theta) d\theta \\ &= \int o(\theta) f_-(\theta) \left. \frac{\partial \epsilon}{\partial R} \right|_{\mu R \text{ fixed}} \frac{d\theta}{2\pi}, \end{aligned} \quad (3)$$

with $f_- = \frac{1}{1 + e^{\epsilon - \mu R}}$. I can now solve the TBA equation iteratively to obtain the expansion

$$\begin{aligned} \langle O(x) \rangle &= \int o(\theta) f_-(\theta) m \cosh \theta \frac{d\theta}{2\pi} + \int o(\theta) f_-(\theta) \frac{d\theta}{2\pi} \int \Phi(\theta - \theta') f_-(\theta') m \cosh \theta' \frac{d\theta'}{2\pi} \\ &\quad + \int o(\theta) f_-(\theta) \frac{d\theta}{2\pi} \int \Phi(\theta - \theta') f_-(\theta') \frac{d\theta'}{2\pi} \int \Phi(\theta' - \theta'') f_-(\theta'') m \cosh \theta'' \frac{d\theta''}{2\pi} + \dots \end{aligned} \quad (4)$$

Notice that this expression in terms of o , Φ and the filling fraction f_- is completely general. It would hold for a system with more thermodynamic couplings, provided these couplings involve conserved quantities. The expression (4) also generalizes easily to cases with several types of particles.

I will now (4) compare with the formula proposed in [1]:

$$\langle O(x) \rangle = \sum_{n=0}^{\infty} \frac{1}{n!} \int \frac{d\theta_1}{2\pi} \cdots \frac{d\theta_n}{2\pi} \prod_{i=1}^n f_-(\theta_i) \langle \theta_n \dots \theta_1 | O(x) \theta_1 \dots \theta_n \rangle_{conn}, \quad (5)$$

and argue that it is the same. Of course, in [1], formula (5) is written only in the case of finite temperature. However, all arguments presented there rely on cancellations between numerators and denominators, together with hints from the free case and the one point function of the stress energy tensor, and these are all features which generalize to more complicated thermodynamic averages, like the ones involving a chemical potential. In (5), the connected form factor is obtained in principle by using the crossing formula for non coinciding arguments, and then getting rid of all the diverging or ill defined terms as the arguments are sent to one another.

The key point here is that the values of the connected form-factors can be understood generally as follows (this argument is quite similar to Balog's construction [5]). To start, I need to make some remarks on the normalization of states. Consider the scalar product $\langle \theta | \theta \rangle$, which, formally, is equal to $2\pi\delta(0)$, since the normalization used in the construction of the asymptotic states is $\langle \theta | \theta' \rangle = 2\pi\delta(\theta - \theta')$. Of course the symbol $\delta(0)$ does not make much sense, and has to be regularized properly. The way to do this is to consider the completeness relation, $1 = \int \frac{d\theta}{2\pi} |\theta\rangle \langle \theta|$. This relation hides the cancellation of two terms, the meaning of which is easier to see by introducing a finite length L in the system. First, the state $|\theta\rangle$ is actually not normalized, because of the $\delta(0)$ term mentioned just before; the integral should therefore involve instead the ket $\frac{|\theta\rangle}{\sqrt{\langle \theta | \theta \rangle}}$ and similarly for the associated bra. Second, allowed rapidities for a particle are *not* uniformly distributed; rather, they have a density given by $2\pi P = mL \cosh \theta$, and the completeness relation should involve therefore an integral $\int P(\theta) d\theta$. For the completeness relation to be equivalent to the one we used so far, which preserves the scalar product of states, we thus need to have [6]

$$\langle \theta | \theta \rangle = mL \cosh \theta. \quad (6)$$

The argument can be generalized to the case with several rapidities. For two particles, the Bethe equations read:

$$\begin{aligned} 2\pi n_1 &= mL \sinh \theta_1 + \frac{1}{i} \ln S(\theta_1 - \theta_2) \\ 2\pi n_2 &= mL \sinh \theta_2 + \frac{1}{i} \ln S(\theta_2 - \theta_1). \end{aligned} \quad (7)$$

From this, we deduce that

$$\begin{aligned} \langle \theta_2, \theta_1 | \theta_1, \theta_2 \rangle &= \text{Det} \begin{vmatrix} mL \cosh \theta_1 + \Phi(\theta_{12}) & -\Phi(\theta_{12}) \\ -\Phi(\theta_{12}) & mL \cosh \theta_2 + \Phi(\theta_{12}) \end{vmatrix} \\ &= m^2 L^2 \cosh \theta_1 \cosh \theta_2 + mL \Phi(\theta_{12}) (\cosh \theta_1 + \cosh \theta_2). \end{aligned} \quad (8)$$

The formula generalizes to an arbitrary number of rapidities. As could have been expected, it coincides with the well known formula for the norm of Bethe states derived by Gaudin and by Korepin in the context of the XXZ and other models (see [7]).

I can now get back to (4) and (5). It is clear that the first term of the expansions match, due to (6)

$$\langle \theta | O(x) | \theta \rangle \equiv \langle \theta | O(x) | \theta \rangle_{conn} = o(\theta) m \cosh \theta. \quad (9)$$

For the two particle form factor, let me define:

$$\begin{aligned} \langle \theta_2, \theta_1 | O(x) | \theta_1, \theta_2 \rangle_{conn} &= \langle \theta_2, \theta_1 | O(x) | \theta_1, \theta_2 \rangle - \langle \theta_2 | O(x) | \theta_2 \rangle_{conn} \langle \theta_1 | \theta_1 \rangle \\ &\quad - \langle \theta_1 | O(x) | \theta_1 \rangle_{conn} \langle \theta_2 | \theta_2 \rangle. \end{aligned} \quad (10)$$

Here, the scalar products (eg $\langle \theta_1 | \theta_1 \rangle$) have to be computed in the presence of the other particle (θ_2), something I left implicit in the notations for simplicity. From the general arguments explained previously, $\langle \theta_1 | \theta_1 \rangle = mL \cosh \theta_1 + \phi(\theta_{12})$. It thus follows that

$$\langle \theta_2, \theta_1 | O(x) | \theta_1, \theta_2 \rangle_{conn} = m \Phi(\theta_{12}) [\cosh \theta_1 o(\theta_2) + \cosh \theta_2 o(\theta_1)], \quad (11)$$

again ensuring a matching between (4) and (5). A general formula for connected form-factors follows easily by extending (10) from $n = 2$ to arbitrary values of n : simply subtract from the initial form factor all the possible contractions, being careful to evaluate scalar products of states in the presence of all other particles - they are all expressed as various minors of the same initial determinant. The net result is simply:

$$\langle \theta_n, \dots, \theta_1 | O(x) | \theta_1, \dots, \theta_n \rangle_{conn} = m \Phi(\theta_{12}) \Phi(\theta_{23}) \dots \Phi(\theta_{n-1,n}) \cosh \theta_1 o(\theta_n) + \text{permutations}, \quad (12)$$

and clearly ensures the coincidence of (4) and (5).

I am of course not pretending that this argument is a proof of (5), but I believe it could be sufficiently strengthened, maybe along the lines of [5]. As it stands, it certainly gives further support to (5), and to my claim that the guess of [1] should be considered for more general thermodynamic ensembles.

I shall now argue that, unfortunately, the formula proposed in [1] for the two particle correlator is probably not correct. Rather than give general arguments, I will present a simple counterexample, in the case where $T = 0$ but there is a non trivial chemical potential.

Consider indeed the free boson with hamiltonian

$$H = \frac{1}{2} \int \left[8\pi g \Pi^2 + \frac{1}{8\pi g} (\partial_x \phi)^2 \right] dx. \quad (13)$$

I will add to this hamiltonian a term $\frac{V}{4\pi} \int \partial_x \phi dx$. Properties in the presence of this coupling are straightforward to evaluate by a simple shift of the bosonic field. Introducing chiral components $\varphi, \bar{\varphi}$, one finds ($\partial_x \phi = \partial_z \varphi + \partial_{\bar{z}} \bar{\varphi}$)

$$\begin{aligned} \langle \partial_z \varphi \rangle &= Vg \\ \langle \partial_{z_1} \varphi \partial_{z_2} \varphi \rangle &= (Vg)^2 - \frac{2g}{(z_1 - z_2)^2}, \end{aligned} \quad (14)$$

where I have set $z = x + iy$, y the imaginary time (called t in [1]).

On the other hand, we can consider the free boson as the UV limit of the sine-Gordon theory, where the hamiltonian (13) is supplemented by an interaction term $\lambda \int \cos \phi dx$. Provided we consider physics at a scale much smaller than the correlation length induced by this perturbation, $\frac{1}{M} \propto \lambda^{-2+2g}$, we will observe results similar to the free boson case. Equivalently, we expect to be able describe the properties of the free boson theory using a massless scattering description [8], with purely right and left moving particles obtained by taking the large rapidity limit of the usual solitons, antisolitons and breathers of the theory. The parametrization of the energy I will use in that limit is $e = \pm p = e^\theta$, where I have set an arbitrary mass scale equal to unity. The correlators (14) should therefore be obtainable using the formulas proposed in [1]. I will show that, actually, only the one point function is obtained.

For simplicity, I restrict to the case $T = 0$. The field V then leads to the creation of a pair of Fermi seas of left and right moving massless solitons: in the following, I will

concentrate on the right moving sector only. The average of $\partial_z \varphi$ is directly related with the number of solitons in the ground state

$$\frac{1}{2\pi} \langle \int \partial_z \varphi dx \rangle = \langle N \rangle$$

This quantity fits exactly in the context described previously. The two particle kernel is given by the soliton soliton S matrix, and, since we are at temperature $T = 0$, the function f_- becomes very simple: $f_- = 1$ for particles in the sea, $f_- = 0$ otherwise. If we consider the operator $\partial_z \varphi$, it couples only to right moving particles, for which the sea is $\theta \in (-\infty, A)$, A a Fermi rapidity. The average of $\partial_z \varphi$ follows from the appropriate generalization of (4):

$$\langle \partial_z \varphi \rangle = \int_{-\infty}^A 2\pi e^\theta \frac{d\theta}{2\pi} + \int_{-\infty}^A 2\pi \frac{d\theta}{2\pi} \int_{-\infty}^A \Phi(\theta - \theta') e^{\theta'} \frac{d\theta'}{2\pi} + \dots \quad (15)$$

Expressions for the form-factors of the operator $\partial_z \varphi$ are well known: we can, in principle, directly compute the connected ones, and show agreement with the general formula (12). For instance, the two particle form-factor gives

$$\langle \theta | \partial_z \varphi | \theta \rangle = 2\pi e^\theta, \quad (16)$$

in agreement with (the massless limit of) (9) for $o(\theta) = 2\pi$, the value of the integral of $\partial_z \varphi$ for a soliton. I have checked similarly the formula for the connected four particle form-factor (11) for simple values of the coupling g .

I now turn to the two point function of the operator $\partial_z \varphi$, and show that the expression proposed for it in [1] this time does not work. Recall the proposal of [1]:

$$\begin{aligned} \langle O(x, y) O(0, 0) \rangle_R &= (\langle O \rangle_R)^2 + \sum_{N=1}^{\infty} \frac{1}{N!} \sum_{\sigma_i = \pm 1} \int \frac{d\theta_1}{2\pi} \dots \frac{d\theta_N}{2\pi} \prod_{j=1}^N f_{\sigma_j}(\theta_j) \\ &\quad \exp \left(i\sigma_j (x\tilde{k}_j + iy\tilde{e}_j) \right) |\langle 0 | O(0) | \theta_1, \dots, \theta_N \rangle_{\sigma_1, \dots, \sigma_N}|^2. \end{aligned} \quad (17)$$

Here, $f_+ = \frac{1}{1+e^{-\epsilon}}$, $\tilde{e} = R\epsilon$ is the dressed excitation energy, while $\tilde{p}$ is the dressed excitation momentum. Finally,

$$\langle 0 | O(0) | \theta_1, \dots, \theta_N \rangle_{\sigma_1, \dots, \sigma_N} \equiv \langle 0 | O(0) | \theta_1 - i\pi(\sigma_1 - 1)/2, \dots, \theta_N - i\pi(\sigma_N - 1)/2 \rangle$$

The physical meaning of (17) is more transparent than the formula. Correlations at temperature T should be determined by processes involving excitations over the thermal ground

state, with dressed energy and momentum [2]. These excitations can be of two types, particles or holes (the variable σ in (17)). A sum as (17) is thus expected; the key proposal of [1] is that the form factors for the physical excitations are the same as the bare ones.

In the case we consider, the situation simplifies considerably. First, let me recall the structure of the excitations [9]. It is conveniently represented in the following table:

$$\begin{aligned}
\text{holes} &: \epsilon_+^h(\theta), \theta \leq A \\
\text{solitons} &: \epsilon_+(\theta), \theta \geq A \\
\text{antisolitons} &: \epsilon_-(\theta) \geq (1-g)V, \theta \text{ arbitrary} \\
\text{breathers} &: \epsilon_n(\theta) \geq ngV, \theta \text{ arbitrary}
\end{aligned} \tag{18}$$

The excitation energies ϵ have non trivial expansions given in [9].

The physical processes involved in the two point function of $\partial_z \varphi$ are of three basic types: creation of a pair particle hole (ie take a soliton in the Fermi sea and move it outside); creation of a pair soliton (above the Fermi sea) antisoliton, and creation of a breather. The latter two processes have thresholds. Since the quantities $\tilde{e}$ and $\tilde{p}$ have no singularity, and the bare form-factors do not know anything about these thresholds, it is immediately quite obvious that the proposal of [1] cannot be true: it would lead to singularities in the Fourier spectrum of the correlator, in sharp contrast with the result expected from (14).

Nevertheless, it is probably reasonable to make this counter example more explicit. First, let me show that the formula is in fact right for the free case $g = \frac{1}{2}$. In that case, things simplify for several reasons: there are no breathers, the excitation energies have simple expressions $\epsilon_+^h = \frac{V}{2} - e^\theta$, $\epsilon_+ = e^\theta - \frac{V}{2}$, $\epsilon_- = e^\theta + \frac{V}{2}$, and only the two particle form factor of $\partial_z \varphi$ is non zero, $\langle 0 | \partial_z \varphi | \theta_1 \theta_2 \rangle = 2i\pi e^{\theta_1/2} e^{\theta_2/2}$. The connected term in expression (17) reads then

$$\begin{aligned}
& \int_{-\infty}^A \frac{d\theta}{2\pi} \int_A^\infty \frac{d\theta_2}{2\pi} \left| 2\pi e^{\theta_1/2} e^{\theta_2/2} \right|^2 e^{iz(e^{\theta_2} - e^{\theta_1})} \\
& + \int_{-\infty}^\infty \frac{d\theta_1}{2\pi} \int_A^\infty \frac{d\theta_2}{2\pi} \left| 2i\pi e^{\theta_1/2} e^{\theta_2/2} \right|^2 e^{iz(e^{\theta_1} + e^{\theta_2})} \\
& = -\frac{1}{z^2} [1 - \exp(e^A z)] - \frac{1}{z^2} \exp(e^A z) = -\frac{1}{z^2}
\end{aligned}$$

the expected result (14) for $g = \frac{1}{2}$.

In the general case, I will consider the Fourier transform of the two point function, $S(\omega) = \int \frac{dx}{2\pi} e^{-i\omega x} \langle \partial_z \varphi(x, y) \partial_z \varphi(0) \rangle_{y \rightarrow 0} = 2g\omega$ ($\omega > 0$). I will also restrict to the case of

low frequencies (that is, ω smaller than any of the thresholds), where only the particle hole processes contribute; if formula (17) fails in that case, it will be enough to show it is not correct.

Below the lowest thresholds, $S(\omega)$, according to formula (17), is given by a sum of terms S_n corresponding to processes with n particles and n holes. When $V = 0$ and we know formula (17) is correct, each of the S_n 's is linear in ω , $S_n = c_n \omega$. In that case, there are no thresholds, so the S_n have to be supplemented by the terms corresponding to the other processes, each of which is also proportional to ω . The infinite sum of all these ω terms reproduces the desired behaviour $S(\omega) = 2g\omega$; as checked in [10] the convergence of this sum is in fact very quick for g not too close to 1. When $V \neq 0$, formula (17) predicts correctly (this simply follows from dimensional analysis) that all the S_n 's now have the form $S_n = \omega f_n\left(\frac{\omega}{V}\right)$, with $c_n = f_n(\infty)$. Let us now consider the limit of small frequencies at finite V : in that case, we are instead exploring the behaviour of the functions f_n in the limit of very small argument. Since the contribution at frequency ω is determined by rapidities $\theta_i > A$ of holes and $\theta'_i < A$ of particles such that

$$\omega = \sum_{i=1}^n \epsilon_+^h(\theta_i) + \sum_{i=1}^n \epsilon_+(\theta'_i), \quad (19)$$

clearly, for small ω , the particles and holes have to be closer and closer to the Fermi rapidity. The two particle contribution for instance, which reads in general

$$\int_{-\infty}^A \frac{d\theta_1}{2\pi} \int_A^{\infty} \frac{d\theta'_1}{2\pi} |\langle \theta'_1 | \partial_z \varphi | \theta_1 \rangle|^2 \delta[\omega - \epsilon_+^h(\theta_1) - \epsilon_+(\theta'_1)]$$

becomes in the limit of small ω , $\frac{1}{(2\pi)^2} |\langle A | \partial_z \varphi | A \rangle|^2 \frac{\omega}{|\dot{\epsilon}_+(A) \dot{\epsilon}_+^h(A)|}$ (dots denote derivatives with respect to the rapidity variable). In fact, for the limit $\lim_{\omega \rightarrow 0} \frac{S(\omega)}{\omega}$, this two particle contribution is the *only* one to consider. There are two reasons for that: one is that the next contribution is down by an ω^2 term due to phase space considerations; the other is that in the limit $\omega \rightarrow 0$, the rapidities of the n particles and n holes are all compressed towards A , and form-factors like eg $\langle A, A | \partial_z \varphi | A, A \rangle$ vanish due to the general behaviour under rapidity exchanges ($S(0) = -1$). It follows that, according to [1], one would find

$$\lim_{\omega \rightarrow 0} \frac{S(\omega)}{|\omega|} \stackrel{?}{=} \frac{1}{(2\pi)^2} \frac{|\langle A | \partial \varphi | A \rangle|^2}{|\dot{\epsilon}_+(A)| |\dot{\epsilon}_+^h(A)|}. \quad (20)$$

The right hand side can be evaluated using the detailed results of [9] for the filling of the ground state etc. One finds (for dimensional reasons, this has to be a pure, V independent number)

$$\text{RHS} = \frac{1}{2\pi} \frac{g^{\frac{1-2g}{1-g}}}{1-g} \left(\frac{\Gamma[g/2(1-g)]}{\Gamma[1/2(1-g)]} \right)^2. \quad (21)$$

Except when $g = \frac{1}{2}$, the right hand side is off the exact result (it should be equal to $2g$) by a finite amount, demonstrating that (17) is not correct in that case.

The reader might worry here that the whole argument assumes somehow absolute convergence of the series in (17), and that maybe one would get the correct limit in (20) by first summing over all contributions, then letting $\omega \rightarrow 0$. The answer to this is that I have considered the limit (20) only to make things as clear as possible. We are of course interested not only in this limit, but in the behaviour of $S(\omega)$ on the whole real axis, where we have to recover $S(\omega) = 2g\omega$ for (17) to be correct. As argued above, only the one particle hole contribution has a term linear in ω : other terms start with higher powers in ω , and it is easy to see that only a finite number of them contribute to a given order in ω . The series representing $S(\omega)$ according to (17) is thus of the form $\sum d_k \omega^k$, with all the d_k 's finite (eqn. (21) means that d_1 is not the right one; one can also check that d_3 is not the right one - that is, it does not vanish - etc). For my argument to be spoilt by a convergence problem, one would need this series to diverge and to somewhat “represent” the simple linear term $2g\omega$; besides the fact that I have found no indication of divergence numerically (that is, convergence seems as good as in [10]), I do not think such a scenario is likely at all.

The reader might also be surprised by the fact that (17) is right for $V = 0$ but not $V \neq 0$. The point is once again that (17) leads to a representation of $S(\omega)$ as a sum of terms of the form $\omega f_n(\frac{\omega}{V})$; validity in the case $V = 0$ is a statement about the sum of $f_n(\infty)$'s, while invalidity in the case $V \neq 0$ is a statement about the general shape of the sum of f_n 's at finite argument.

The physical origin of the failure of (17) is easy to trace back to the dressing effects. In fact, in [9], another approach to compute correlators at $T = 0$ with a field coupled to the $U(1)$ current was proposed. In this paper, it is recognized that dressed excitations must have dressed form-factors; an expression was proposed in particular for the low energy behaviour of the particle hole form factor, which reproduced $S(\omega)$ correctly.

I should stress that I have not found simple counterexamples in the context originally considered in [1], of a theory with a temperature and no chemical potential. However, I

believe the argument presented in [1] generalizes to any thermodynamic average, so this counterexample at $T = 0$ with a chemical potential is nevertheless a good, alas negative, test, of their result for two and higher point functions. Some easy things can be said about the $T \neq 0$ case however. One of them has to do with *thresholds*. Consider for instance the massless limit of the sine-Gordon model once again, and its massless scattering description. At $T \neq 0$, it is one of the striking features of the interactions that the energy necessary to add a particle has a gap: for instance, the energy to create a one-breather in the attractive regime is $T\epsilon_1 \geq T \ln 3$; similarly the energy gained by destroying such a breather is larger or equal to this number (this follows simply from the solution of the TBA equations). As a result, if we consider again the correlator of $\partial_z \varphi$, processes involving the one-breather have thresholds (recall that the bare form-factors of $\partial_z \varphi$ with an even number of one-breathers are zero). Since the form factors in the sum (17) are the bare ones, they know nothing about these thresholds; as for the other pseudo energies, they have no singularity at the position of these thresholds. It follows that, according to (17), the Fourier transform of the two point function of $\partial_z \varphi$ would exhibit singularities at finite values of the frequency ω . This is of course in contradiction with the simple form of this two point function, that follows from conformal invariance, and indicates once again that the proposal in [1] is generally not correct.

In conclusion, I believe that the formulas of [1] for the two and higher point functions are in general incorrect because they don't take properly into account the dressing of the effective vacuum created by finite thermodynamic couplings. As for the one point function of conserved quantities, the formula of [1] looks very reasonable when compared with the result of the TBA, and presumably could be rigorously proven by a more serious analysis of connected form-factors than the one I have presented here. I am not sure about the one point functions of non conserved quantities - the examples studied in [1] are quite convincing, but further work is probably needed to settle the issue.

Acknowledgments: I thank A. Leclair, F. Lesage and G. Mussardo for many discussions on this problem. This work was supported by the DOE and the NSF (under the NYI program).

References

- [1] A. Leclair and G. Mussardo, “Finite temperature correlation functions in integrable QFT”, hep-th/9902075.
- [2] A. LeClair, F. Lesage, S. Sachdev, and H. Saleur, Nucl. Phys. B482 (1996) 579.
- [3] S. Sachdev and K. Damle, Phys. Rev. Lett. 78 (1997) 943.
- [4] Al. Zamolodchikov, Nucl. Phys. B342 (1990) 695.
- [5] J. Balog, Nucl. Phys. B419 (1994) 480.
- [6] A. Leclair, G. Mussardo, H. Saleur, and S. Skorik, Nucl. Phys. B453 (1995) 581.
- [7] V.E. Korepin, N.M. Bogoliubov and A.G.Izergin, “Quantum inverse scattering method and correlation functions”, Cambridge University Press, 1993.
- [8] P. Fendley, H. Saleur and N. P. Warner, Nucl. Phys. B430 (1994) 577.
- [9] F. Lesage and H. Saleur, Nucl. Phys. B490 (1997) 543.
- [10] F. Lesage, H. Saleur, and S. Skorik, Nucl. Phys. B474 (1996) 602.